

February 1, 2008

How to build a non-spreading wave packet in quantum mechanics

S. Bruce
Department of Physics
University of Concepcion
P.O. Box 4009
Concepcion, Chile

Abstract

Quantum mechanics asserts that a wave packet must inevitably spread as time progresses since the dispersion relation for the quantum waves is assumed to be quadratic in the momentum k . However, this assumption does not consider the standard frequency Doppler shift formula of Galilean relativity. In this article a non-dispersive wave packet is constructed by appropriately considering the transformation rules between the laboratory and the (particle's) rest inertial reference frames.

PACS 03.65.Bz - Foundations, theory of measurement, miscellaneous theories.

E-mail: sbruce@udec.cl

Fax: (56-41) 22 4520.

As is well known wave packets for particles of nonzero rest mass are dispersed while they evolve in time. This is because wave-like properties must be attributed to the quantum system, the wave vector $\mathbf{k}$ and the frequency ω being related to the momentum and energy by

$$\mathbf{p} \rightarrow \hbar \mathbf{k}, \quad E \rightarrow \hbar \omega(\mathbf{k}). \quad (1)$$

In the classical (non-relativistic) level the dependence of the energy on the momentum must therefore be given by the familiar relationship $E(\mathbf{p}) = \mathbf{p}^2/2m_0$, where m_0 is the rest mass of the particle. The corresponding dispersion relation for the *quantum waves* in the free-particle case is therefore

$$\omega(\mathbf{k}) = \frac{\hbar}{2m_0} \mathbf{k}^2. \quad (2)$$

Consider now a non-relativistic wave packet

$$\Psi(\mathbf{x}, t) = \frac{1}{(2\pi\hbar)^{3/2}} \int d^3k \exp[i\mathbf{k} \cdot \mathbf{x}] \Phi(\mathbf{k}, t). \quad (3)$$

We assume Ψ is confined to a spatial region $V_{\mathbf{x}}$. We may construct such a packet by means of a function $\Phi(\mathbf{k}, t)$ which is concentrated in a volume $V_{\mathbf{k}}$ of k -space whose dimensions are inversely proportional to $V_{\mathbf{x}}$. Given the wave function at some instant t_0 ($= 0$) throughout all the space, we can compute Ψ at any other future time. This follows from (3):

$$\Psi(\mathbf{x}, t) = \frac{1}{(2\pi\hbar)^{3/2}} \int d^3k \exp[i(\mathbf{k} \cdot \mathbf{x} - \omega(\mathbf{k})t)] \Phi(\mathbf{k}, 0). \quad (4)$$

Thus we can write

$$\Psi(\mathbf{x}, t) = \int d^3x' G(\mathbf{x} - \mathbf{x}', t) \Psi(\mathbf{x}', 0), \quad (5)$$

where

$$G(\mathbf{x}, t) \equiv \frac{1}{(2\pi)^{3/2}} \int d^3k \exp[i(\mathbf{k} \cdot \mathbf{x} - \omega(\mathbf{k})t)] = \left(\frac{m_0}{2\pi i \hbar t}\right)^{3/2} \exp\left[\frac{im_0 |\mathbf{x}|^2}{2\hbar t}\right] \quad (6)$$

is the well known *free-particle* Green function.¹ A particular example of considerable interest is the one-dimensional gaussian wave function

$$\Psi(x, 0) = A \exp(-(x - x_0)^2/2a^2) \exp(ip_x x/\hbar). \quad (7)$$

From (5) and (6) we get²

$$\begin{aligned} \Psi(x,t) &= \pi^{-1/4} \left(a + \frac{i\hbar t}{m_0 a} \right)^{-1/2} \exp \left[\frac{ip_x x}{\hbar} - \frac{ip_x^2 t}{2m_0 \hbar} \right] \\ &\times \exp \left[-\frac{(x - x_0 - p_x t/m_0)^2 (1 - i\hbar/m_0 a^2)}{2(a^2 + \hbar^2 t^2/m_0^2 a^2)} \right]. \end{aligned} \quad (8)$$

The corresponding probability distribution is

$$P(x,t) = |\Psi(x,t)|^2 = \pi^{-1/2} \left(a^2 + \frac{\hbar^2 t^2}{m_0^2 a^2} \right)^{-1/2} \exp \left[-\frac{(x - x_0 - p_x t/m_0)^2}{(a^2 + \hbar^2 t^2/m_0^2 a^2)} \right]. \quad (9)$$

Notice that $G(\mathbf{x},t)$ is the solution which evolves out of the Dirac delta function $\delta(\mathbf{x})$ at $t = 0$. However, for $t > 0$, G will make $\Psi(\mathbf{x},t)$ and $P(\mathbf{x},t)$ to spread out in space.

Let K and K' be two inertial reference frames. The K' system has all its axes parallel to those of K and moves with velocity $\mathbf{v}$ relative to K . Assume E, p and E', p' to be the energy and momentum of a particle in the K and K' frames, respectively. The Lorentz transformation for these quantities are

$$E(\mathbf{p}) = \mathbf{v} \cdot \mathbf{p} + \frac{E'}{\gamma}, \quad (10)$$

$$\mathbf{p} = \mathbf{p}' + (\gamma - 1) \frac{\mathbf{p}' \cdot \mathbf{v}}{v^2} \mathbf{v} + \gamma \frac{\mathbf{v}}{c^2} E', \quad (11)$$

where $\gamma = (1 - v^2/c^2)^{-1/2}$. Let us now assume that K' is the rest reference frame of the particle. Equation (10) becomes

$$E(\mathbf{p}) = \mathbf{v} \cdot \mathbf{p} + \frac{m_0 c^2}{\gamma}. \quad (12)$$

It is interesting to note that this expression represents a correspondence relation for the E_D -energy eigenvalues of the Dirac Hamiltonian. We can see this by calculating the energy eigenvalues of the Dirac energy operator $H_D = c\boldsymbol{\alpha} \cdot \mathbf{p} + m_0 c^2 \beta$ in the energy-helicity eigenstates $|E_D, s\rangle$:

$$\begin{aligned} E_D &\equiv \langle E_D, s | H_D | E_D, s \rangle \\ &= \langle E_D, s | c\boldsymbol{\alpha} \cdot \mathbf{p} + m_0 c^2 \beta | E_D, s \rangle \\ &= \langle E_D, s | c\boldsymbol{\alpha} | E_D, s \rangle \cdot \mathbf{p} + \langle E_D, s | \beta | E_D, s \rangle m_0 c^2. \end{aligned} \quad (13)$$

Furthermore

$$\begin{aligned} < E_D, s | c\boldsymbol{\alpha} | E_D, s > = c^2 \frac{\mathbf{p}}{E_D} = \mathbf{v} , \\ < E_D, s | \beta | E_D, s > = 1/\gamma , \end{aligned} \quad (14)$$

as can be easily checked ¹. Then from (12) and (13) we can make the correspondence $E \longleftrightarrow E_D$. It is important to observe that a separation of concepts is involved in (12) and (14). The term $\mathbf{v} \cdot \mathbf{p}$ in (12) is the scalar product of the velocity $\mathbf{v}$ of the frame K' , where the particle is at rest, and the momentum $\mathbf{p}$ of the particle, both relative to the (say, laboratory) frame K . They are velocities associated with different objects. Therefore, at the quantum level, we have to distinguish between $\mathbf{p}/m_0$ and $\mathbf{v}$ although they have the same *mean* values.

For the sake of simplicity, let us now assume the non-relativistic limit of (12). If $|\mathbf{v}/c|^2 \ll 1$, we get

$$E(\mathbf{p}) = \mathbf{v} \cdot \mathbf{p} - \frac{1}{2}m_0v^2 + m_0c^2 . \quad (15)$$

Accordingly, the dispersion relation for the wave function (4) is

$$\omega(\mathbf{k}) = \mathbf{v} \cdot \mathbf{k} - \frac{\omega' v^2}{2 c^2} + \omega', \quad \omega' = \frac{m_0c^2}{\hbar} . \quad (16)$$

From (16) and (4) we can build up a free-particle wave packet that moves with group velocity $\mathbf{v} = \partial\omega(\mathbf{k})/\partial\mathbf{k}$. From (??) and (16) we find

$$\begin{aligned} \Psi(\mathbf{x}, t) &= \frac{1}{(2\pi\hbar)^{3/2}} \int d^3k \exp \left[i\mathbf{k} \cdot (\mathbf{x} - \mathbf{v}t) + i\left(\frac{\omega' v^2}{2 c^2} - \omega'\right)t \right] \Phi(\mathbf{k}, 0) \\ &= \exp \left[i \left(\frac{\omega' v^2}{2 c^2} - \omega' \right) t \right] \frac{1}{(2\pi\hbar)^{3/2}} \int d^3k \exp [i\mathbf{k} \cdot (\mathbf{x} - \mathbf{v}t)] \Phi(\mathbf{k}, 0) \\ &= \exp \left[i \left[\frac{\omega' v^2}{2 c^2} - \omega' \right] t \right] \Psi(\mathbf{x} - \mathbf{v}t, 0) . \end{aligned} \quad (17)$$

Therefore, we can write

$$\Psi(\mathbf{x}, t) = \int d^3x' \mathcal{G}(\mathbf{x} - \mathbf{x}', t) \Psi(\mathbf{x}', 0) , \quad (18)$$

¹Here $\boldsymbol{\alpha}$ and β are the usual Dirac matrices.

with

$$\mathcal{G}(\mathbf{x}, t) = \exp \left[-\frac{im_0c^2}{\hbar} \left(1 - \frac{v^2}{c^2} \right) t \right] \delta^{(3)}(\mathbf{x} - \mathbf{v}t) , \quad (19)$$

the corresponding Green function. This shows that the wave packet (17) evolves without deformation with velocity $\mathbf{v}$ relative to the frame K .

Note that (16) considers the Doppler effect of the quantum waves as the result of the movement of the wave packet relative to an observer in K . The consequence is the outcome of an unspreading wave packet. Correspondingly, the probability density is a function that progresses without deformation:

$$P(\mathbf{x}, t) = | \Psi(\mathbf{x} - \mathbf{v}t, 0) |^2 . \quad (20)$$

This work was supported by Dirección de Investigación, Universidad de Concepción, through grant P.I. 96.11.19-1.0. and by Fondecyt grant # 1970995.

References

¹K. Gottfried, *Quantum Mechanics* Vol 1 (W. A. Benjamin, 1966).

²R. B. Leighton, *Principles of Modern Physics* (McGraw-Hill NY, 1959).